



Mark Scheme (Results)

January 2024

Pearson Edexcel International Advanced
Subsidiary Level in Chemistry (WCH12)
Paper 01 Energetics, Group Chemistry,
Halogenoalkanes and Alcohols

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2024

Question Paper Log Number P75592A

Publications Code WCH12_01_2401_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.
() means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer. ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities. Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Section A

Question Number	Answer	Mark
1	The only correct answer is A (HF) <i>B is not correct because tellurium and hydrogen have the same electronegativity</i> <i>C is not correct because arsenic is less electronegative than hydrogen</i> <i>D is not correct because tin is less electronegative than hydrogen</i>	(1)
2	The only correct answer is B (H₂O(l) → H₂O(g)) <i>A is not correct because a covalent bond is being broken</i> <i>C is not correct because only London forces are being broken</i> <i>D is not correct because covalent bonds are being broken and formed</i>	(1)
3	The only correct answer is C (PH₃) <i>A is not correct because NO has polar bonds</i> <i>B is not correct because BeCl₂ has polar bonds</i> <i>D is not correct because CCl₄ is symmetrical and has only London forces between its molecules</i>	(1)

Question Number	Answer	Mark
4	<p>The only correct answer is D ($\text{CH}_3(\text{CH}_2)_7\text{CH}_3$)</p> <p><i>A is not correct because $(\text{CH}_3)_4\text{C}$ has fewer electrons and a branched carbon chain</i></p> <p><i>B is not correct because $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ has fewer electrons</i></p> <p><i>C is not correct because $(\text{C}_2\text{H}_5)_4\text{C}$ has a branched carbon chain</i></p>	(1)

Question Number	Answer	Mark
5	<p>The only correct answer is B (cyclohexane)</p> <p><i>A is not correct because ammonia is a polar solvent</i></p> <p><i>C is not correct because methanol is a polar solvent</i></p> <p><i>D is not correct because water is a polar solvent</i></p>	(1)

Question Number	Answer	Mark
6	<p>The only correct answer is D (CH_4)</p> <p><i>A is not correct because the oxidation state of F is -1 in HF</i></p> <p><i>B is not correct because the oxidation state of H is -1 in NaH</i></p> <p><i>C is not correct because the oxidation state of O is -1 in H_2O_2</i></p>	(1)

Question Number	Answer	Mark
7	<p>The only correct answer is A ($\text{Br}_2 + \text{F}_2 \rightarrow 2\text{BrF}$)</p> <p><i>B is not correct because chlorine is reduced and fluorine does not change oxidation state</i></p> <p><i>C is not correct because fluorine is reduced only</i></p> <p><i>D is not correct because this is not a redox reaction</i></p>	(1)

Question Number	Answer	Mark
8	<p>The only correct answer is B (BaCl_2 and Ag_2SO_4)</p> <p><i>A is not correct because in B $\text{BaSO}_4(s)$ is formed as well as $\text{AgCl}(s)$</i></p> <p><i>C is not correct because no solid would form</i></p> <p><i>D is not correct because in B $\text{AgCl}(s)$ is formed as well as $\text{BaSO}_4(s)$</i></p>	(1)

Question Number	Answer	Mark
9	<p>The only correct answer is C (nitric acid)</p> <p><i>A is not correct because sodium carbonate is used to identify carboxylic acids</i></p> <p><i>B is not correct because sodium hydroxide is used to identify ammonium ions</i></p> <p><i>D is not correct because ammonia is used to distinguish silver halide precipitates</i></p>	(1)

Question Number	Answer	Mark
10	<p>The only correct answer is D (40)</p> <p><i>A is not correct because a mole ratio of 2:1 instead of 1:2 has been used and the volume is in dm³</i></p> <p><i>B is not correct because this is the correct volume in dm³</i></p> <p><i>C is not correct because a mole ratio of 2:1 instead of 1:2 has been used</i></p>	(1)

Question Number	Answer	Mark
11	<p>The only correct answer is C (2.8 g)</p> <p><i>A is not correct because this is the percentage by mass of nitrogen in ammonium nitrate</i></p> <p><i>B is not correct because this is the mass of ammonium nitrate</i></p> <p><i>D is not correct because only one nitrogen atom has been taken into account</i></p>	(1)

Question Number	Answer	Mark
12	<p>The only correct answer is A (increases, decreases, decreases)</p> <p><i>B is not correct because the boiling temperature increases, the electronegativity decreases and the reactivity as an oxidising agent decreases down the group</i></p> <p><i>C is not correct because the reactivity as an oxidising agent decreases down the group</i></p> <p><i>D is not correct because the boiling temperature increases and the electronegativity decreases down the group</i></p>	(1)

Question Number	Answer	Mark
13	<p>The only correct answer is D (the frequency of collisions with $E \geq E_a$ increases)</p> <p><i>A is not correct because the activation energy remains the same</i></p> <p><i>B is not correct because there is no reference to the frequency or energy of collisions</i></p> <p><i>C is not correct because the particles must collide in order to react</i></p>	(1)

Question Number	Answer	Mark
14	<p>The only correct answer is A (the position of equilibrium is affected by temperature, by pressure and by catalysts)</p> <p><i>B is not correct because the concentration of the reactants remain constant</i></p> <p><i>C is not correct because the rate of the forward reaction is equal to the rate of the backward reaction</i></p> <p><i>D is not correct because equilibrium can be reached from either direction</i></p>	(1)

Question Number	Answer	Mark
15	<p>The only correct answer is D (1-iodobutane)</p> <p><i>A is not correct because the carbon-halogen bond enthalpy decreases down the group</i></p> <p><i>B is not correct because the carbon-halogen bond enthalpy decreases down the group</i></p> <p><i>C is not correct because the carbon-halogen bond enthalpy decreases down the group</i></p>	(1)

Question Number	Answer	Mark
16	<p>The only correct answer is C (4)</p> <p><i>A is not correct because there are 4 cyclic alcohol structural isomers with molecular formula C₄H₈O</i></p> <p><i>B is not correct because there are 4 cyclic alcohol structural isomers with molecular formula C₄H₈O</i></p> <p><i>D is not correct because there are 4 cyclic alcohol structural isomers with molecular formula C₄H₈O</i></p>	(1)

Question Number	Answer	Mark
17	<p>The only correct answer is C (CH₃CH₂CH₂CH₂OH + 4O₂ → 4CO + 5H₂O)</p> <p><i>A is not correct because this equation shows complete combustion</i></p> <p><i>B is not correct because this equation shows complete combustion and the oxygen is not balanced</i></p> <p><i>D is not correct because the oxygen is not balanced</i></p>	(1)

Question Number	Answer	Mark
18	<p>The only correct answer is A (1, 2 and 3)</p> <p><i>B is not correct because tertiary alcohols react with concentrated HCl to produce chloroalkanes</i></p> <p><i>C is not correct because alcohols react with PCl₅ to produce chloroalkanes</i></p> <p><i>D is not correct because concentrated H₂SO₄ and KCl produce HCl which reacts with the tertiary alcohol</i></p>	(1)

Question Number	Answer	Mark
19	<p>The only correct answer is D (5)</p> <p><i>A is not correct because each primary alcohol group needs 2 mol of [O] and the secondary alcohol group needs 1 mol of [O]</i></p> <p><i>B is not correct because each primary alcohol group needs 2 mol of [O] and the secondary alcohol group needs 1 mol of [O]</i></p> <p><i>C is not correct because each primary alcohol group needs 2 mol of [O] and the secondary alcohol group needs 1 mol of [O]</i></p>	(1)

Question Number	Answer	Mark
20	<p>The only correct answer is B (CH₃COCH₃)</p> <p><i>A is not correct because the molecular ion peak would be at m/z = 60 and there would be a fragment ion peak at m/z = 45</i></p> <p><i>C is not correct because there would be a fragment ion peak at m/z = 29</i></p> <p><i>D is not correct because there would be a fragment ion peak at m/z = 29</i></p>	(1)

(Total for Section A = 20 marks)

Section B

Question Number	Answer	Additional Guidance	Mark
21(a)	An answer that makes reference to the following point: <ul style="list-style-type: none"><li data-bbox="353 443 689 480">• $1s^22s^22p^63s^23p^64s^2$	Accept correct electrons-in-boxes notation Accept expansion of p-subshell(s), eg $2p_x^22p_y^22p_z^2$ for $2p^6$ Allow non-superscript numbers for electrons Ignore use of commas and spaces Ignore 2,8,8,2 Ignore [Ar] for $1s^22s^22p^63s^23p^6$ Do not award incorrect order of subshells	(1)

Question Number	Answer	Additional Guidance	Mark
21(b)(i)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> calcium/Ca oxidised and from 0 (in Ca) to (+)2 (in CaCl₂) (1) chlorine/Cl₂/Cl reduced and from 0 (in Cl₂) to -1 (in CaCl₂) (1) 	<p>Ignore explanations in terms of electron loss/gain</p> <p>Oxidation numbers may be shown in the equation</p> <p>Allow calcium/Ca is a reducing agent</p> <p>Ignore just oxidation number of Ca increases</p> <p>Allow chlorine/Cl₂/Cl is an oxidising agent</p> <p>Ignore just oxidation number of Cl decreases</p> <p>If no other mark awarded: Calcium/Ca changes from 0 to +2 and chlorine/Cl₂/Cl changes from 0 to -1 scores (1) OR Calcium/Ca oxidised/reducing agent and chlorine/Cl₂/Cl reduced/oxidising agent scores (1) provided no contradiction in terms of increase/decrease in oxidation number</p>	(2)

Question Number	Answer	Additional Guidance	Mark												
21(b)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> any two or three correct scores (1) any four or five correct scores (2) all six correct scores (3) 	<p><u>Example of completed table:</u></p> <table border="1"> <thead> <tr> <th>Species</th> <th>Bonding</th> <th>Structure</th> </tr> </thead> <tbody> <tr> <td>(Ca(s))</td> <td>metallic</td> <td>giant / lattice</td> </tr> <tr> <td>Cl₂((g))</td> <td>(covalent)</td> <td>simple / molecular</td> </tr> <tr> <td>CaCl₂((s))</td> <td>ionic</td> <td>(giant)</td> </tr> </tbody> </table>	Species	Bonding	Structure	(Ca(s))	metallic	giant / lattice	Cl ₂ ((g))	(covalent)	simple / molecular	CaCl ₂ ((s))	ionic	(giant)	(3)
Species	Bonding	Structure													
(Ca(s))	metallic	giant / lattice													
Cl ₂ ((g))	(covalent)	simple / molecular													
CaCl ₂ ((s))	ionic	(giant)													

Question Number	Answer	Additional Guidance	Mark
21(b)(iii)	An answer that makes reference to the following point: <ul style="list-style-type: none"><li data-bbox="376 373 595 408">• orange-red	Accept brick-red Allow red Allow orange Allow yellow-red Allow yellow-orange Ignore just yellow Do not award crimson red Do not award scarlet red Do not award magenta red	(1)

Question Number	Answer	Additional Guidance	Mark
21(c)(i)	<p>An answer that makes reference to any two of the following points:</p> <ul style="list-style-type: none"> <li data-bbox="360 373 1070 408">• P1: brown gas (produced) (1) <li data-bbox="360 624 1070 659">• P2: bubbles (through water) (1) <li data-bbox="360 767 1070 802">• P3: (water/indicator) turns red (1) <li data-bbox="360 1018 1070 1053">• P4: solid melts (1) 	<p>Penalise incorrect gas, including other nitrogen oxides (eg NO), in P1/P2 once only</p> <p>Allow fumes for gas Ignore red Do not award any other colour Ignore nitrogen dioxide/NO₂ Ignore oxygen/O₂ Ignore any reference to relighting of a glowing splint</p> <p>Ignore effervescence/fizzing Ignore nitrogen dioxide/NO₂/oxygen/O₂ Ignore air/nitrogen/N₂</p> <p>Allow turns pink Ignore turns orange Ignore turns yellow Do not award any other colour Do not award water turns milky / white ppt formed If stated, initial colour must be green/yellow</p> <p>Allow colourless liquid formed Ignore solid glows Ignore white solid/ppt produced (in heated tube) Do not award solid dissolves Do not award black solid produced</p>	(2)

Question Number	Answer	Additional Guidance	Mark
21(c)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • magnesium (nitrate) is less (thermally) stable (1) • magnesium ion/Mg²⁺ is smaller OR magnesium ion/Mg²⁺ has higher charge density and is more polarising (1) • less energy needed to break N–O (1) 	<p>Accept reverse arguments throughout</p> <p>Allow (thermal) stability increases down the group</p> <p>Allow decomposition/reaction is less endothermic with magnesium (nitrate)</p> <p>Allow decomposition/reaction occurs at a lower temperature with magnesium (nitrate)</p> <p>Allow magnesium (nitrate) decomposes faster / more easily</p> <p>Ignore just magnesium (nitrate) reacts faster / more easily</p> <p>Ignore magnesium ion for magnesium (nitrate)</p> <p>Allow magnesium/Mg has smaller ionic radius</p> <p>Allow magnesium ion/Mg²⁺ has fewer shells</p> <p>Ignore any reference to just magnesium/Mg (atoms)</p> <p>Allow distorts/weakens nitrate/anion more</p> <p>Allow less energy needed to break bond(s) in nitrate/anion</p> <p>Allow easier to break N–O / bond(s) in nitrate/anion</p> <p>Ignore just weaker N–O / bond(s) in nitrate/anion</p> <p>Do not award less energy needed to break (ionic) bond(s) between Mg²⁺/cations and NO₃⁻/anions</p>	(3)

Question Number	Answer	Additional Guidance	Mark																				
*21(d)	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="320 587 1153 882"> <thead> <tr> <th>Number of indicative marking points seen in answer</th> <th>Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td>6</td> <td>4</td> </tr> <tr> <td>5-4</td> <td>3</td> </tr> <tr> <td>3-2</td> <td>2</td> </tr> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> <table border="1" data-bbox="320 991 1153 1458"> <thead> <tr> <th></th> <th>Number of marks awarded for structure and lines of reasoning</th> </tr> </thead> <tbody> <tr> <td>Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning throughout.</td> <td>2</td> </tr> <tr> <td>Answer is partially structured with some linkages and lines of reasoning.</td> <td>1</td> </tr> <tr> <td>Answer has no linkages between points and is unstructured.</td> <td>0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0		Number of marks awarded for structure and lines of reasoning	Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning throughout.	2	Answer is partially structured with some linkages and lines of reasoning.	1	Answer has no linkages between points and is unstructured.	0	<p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> <p>If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded, do not deduct mark(s).</p> <p>Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points																						
6	4																						
5-4	3																						
3-2	2																						
1	1																						
0	0																						
	Number of marks awarded for structure and lines of reasoning																						
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning throughout.	2																						
Answer is partially structured with some linkages and lines of reasoning.	1																						
Answer has no linkages between points and is unstructured.	0																						

<p>Indicative points:</p> <ul style="list-style-type: none"> • IP1: $\text{Ca}(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{Ca}(\text{OH})_2(\text{aq}) + \text{H}_2(\text{g})$ • IP2: effervescence • IP3: (indicator) turns blue • IP4: faster / more vigorous reaction with calcium and linked to an observation <p>OR</p> <p>no (observable) reaction with magnesium</p> <ul style="list-style-type: none"> • IP5: Ca has lower ionisation energy OR Ca has more shielding / more shells / larger atoms • IP6: darker blue / more alkaline / higher pH with Ca and linked to greater solubility of $\text{Ca}(\text{OH})_2$ 	<p>Ignore any reference to Ca/Mg dissolving</p> <p>Allow multiples Ignore state symbols Ignore equations involving magnesium</p> <p>Allow fizzing/bubbling</p> <p>Allow turns purple If stated, initial colour must be green/yellow Ignore forms solution with $\text{pH} > 7$</p> <p>Accept reverse arguments eg reaction with calcium:</p> <ul style="list-style-type: none"> • has faster bubbling • changes colour faster • forms (white) ppt faster • gets hotter (Ignore more exothermic) • (solid) disappears faster <p>Ignore (indicator) remains green with Mg Ignore no (white) ppt with Mg</p> <p>Accept reverse arguments Allow Ca has less attraction on (outer) e^{-} Allow Ca loses electron(s) more easily Ignore calcium ions/Ca^{2+} for calcium/Ca</p> <p>Accept reverse arguments Ignore less (white) ppt with Ca</p> <p>Allow hydroxide solubility increases down group Allow $\text{Mg}(\text{OH})_2$ insoluble Ignore explanations of solubility</p>	
--	--	--

(Total for Question 21 = 18 marks)

Question Number	Answer	Additional Guidance	Mark
22(a)	An explanation that makes reference to the following points: <ul style="list-style-type: none"> • 1 mol of substance / compound / MgO (1) • formed from element(s) in standard state(s) (1) 	Allow 1 mol of product Ignore Mg(s) and O ₂ (g) / reactant(s) for element(s) Ignore normal/natural etc for standard Ignore any reference to standard conditions	(2)


Question Number	Answer	Additional Guidance	Mark
22(b)	<ul style="list-style-type: none"> <li data-bbox="376 804 1014 836">• calculation of energy transferred (1) <li data-bbox="376 948 1014 979">• calculation of amount of MgO (1) <li data-bbox="376 1091 1014 1123">• calculation of $\Delta_r H_2$ (1) <li data-bbox="376 1305 1014 1369">• negative sign and answer to 3SF or 2SF (1) 	<p data-bbox="1055 304 1525 336">Here and throughout the paper:</p> <ul style="list-style-type: none"> <li data-bbox="1099 341 1928 405">• penalise incorrect rounding once only and only if the final answer is incorrect <li data-bbox="1099 410 1794 442">• do not penalise correct premature rounding <li data-bbox="1099 446 1644 478">• penalise incorrect units once only <li data-bbox="1099 483 1442 515">• Allow mol⁻ for mol⁻¹ <p data-bbox="1055 555 1397 587"><u>Example of calculation:</u></p> <p data-bbox="1055 592 1951 624">Correct answer to 3SF or 2SF with some working scores (4)</p> <p data-bbox="1055 663 1883 695">Ignore SF except 1SF and penalise use of 1SF once only</p> <p data-bbox="1055 735 1435 767">Ignore sign in M1 and M3</p> <p data-bbox="1055 807 1621 903">energy = $25.0 \times 4.18 \times (28.0 - 21.5)$ = 679.25 (J) Allow 0.67925 (kJ)</p> <p data-bbox="1055 951 1794 1046">amount = $0.189 \div 40.3$ = $0.0046898 / 4.6898 \times 10^{-3}$ (mol) Allow $0.004725 / 4.725 \times 10^{-3}$ (mol) from $M_r = 40$</p> <p data-bbox="1055 1094 1883 1270">$\Delta_r H_2 = 679.25 \div 0.0046898$ = 144830 (J mol⁻¹) Accept 144.83 (kJ mol⁻¹) Allow 143760 (J mol⁻¹) / 143.76 (kJ mol⁻¹) from $M_r = 40$ TE on M1 and M2</p> <p data-bbox="1055 1310 1845 1445">-145 / -140 (kJ mol⁻¹) Allow -145000 / -140000 J mol⁻¹ Allow -144 (kJ mol⁻¹) / -144000 J mol⁻¹ from $M_r = 40$ TE on M3</p>	(4)

Question Number	Answer	Additional Guidance	Mark
22(c)	<ul style="list-style-type: none"> • correct expression for $\Delta_r H_1$ (1) • calculation of $\Delta_r H_1$ (1) 	<p><u>Example of calculation:</u></p> <p>Correct answer with some working scores (2)</p> <p>$\Delta_r H_1 = -462 + (-286) - \text{answer to (b)}$</p> <p>$\Delta_r H_1 = -462 + (-286) - (-145)$ $= -603 \text{ (kJ mol}^{-1}\text{)}$</p> <p>Ignore SF TE on answer to (b) No TE on incorrect expression from M1</p> <p>Allow $-604 \text{ (kJ mol}^{-1}\text{)}$ from $M_r = 40$ in (b)</p> <p>If using $-100 \text{ (kJ mol}^{-1}\text{)}$ for answer to (b) $\Delta_r H_1 = -462 + (-286) - (-100)$ $= -648 \text{ (kJ mol}^{-1}\text{)}$</p>	(2)

(Total for Question 22 = 8 marks)


Question Number	Answer	Additional Guidance	Mark
23(a)(i)	An answer that makes reference to the following point: <ul style="list-style-type: none"> 2-bromo-2-methylpropane 	Ignore omission of hyphens or use of commas Allow 2-methyl-2-bromopropane Do not award 2,2-bromomethylpropane	(1)

Question Number	Answer	Additional Guidance	Mark
23(a)(ii)	An answer that makes reference to the following point: <ul style="list-style-type: none"> displayed formula of 2-bromobutane 	<u>Example of displayed formula:</u> $ \begin{array}{ccccccc} & \text{H} & \text{Br} & \text{H} & \text{H} & & \\ & & & & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{H} & \\ & & & & & & \\ & \text{H} & \text{H} & \text{H} & \text{H} & & \end{array} $ Ignore any other type of formula Ignore bond lengths and bond angles Ignore any name even if incorrect Do not award non-displayed CH ₃ groups Do not award any missing hydrogens / bonds	(1)

Question Number	Answer	Additional Guidance	Mark
23(a)(iii)	An answer that makes reference to the following point: <ul style="list-style-type: none"> skeletal formula of 1-bromo-2-methylpropane 	<u>Example of skeletal formula:</u>  Ignore any other type of formula Ignore bond lengths and bond angles Ignore any name even if incorrect	(1)

Question Number	Answer	Additional Guidance	Mark
23(b)	An answer that makes reference to the following point: <ul style="list-style-type: none"><li data-bbox="360 373 741 405">• structure of (CH₃)₃CBr	Allow skeletal, structural or displayed formulae, or any combination of these If more than one type of formula used, all must be correct Ignore CH ₃ -C connectivity Ignore name even if incorrect	(1)

Question Number	Answer	Additional Guidance	Mark																				
23(c)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • any two or three correct scores (1) • any four or five correct scores (2) • any six or seven correct scores (3) • all eight correct scores (4) 	<p><u>Example of completed table:</u></p> <p>Allow indication of conditions in reagent box, eg NaOH(aq) indicates aqueous conditions</p> <p>Penalise ions for reagents and/or names for formulae once only</p> <table border="1" data-bbox="996 550 1928 1401"> <thead> <tr> <th data-bbox="996 550 1151 657">Reaction</th> <th data-bbox="1151 550 1373 657">Name of mechanism</th> <th data-bbox="1373 550 1568 657">Formula of Reagent</th> <th data-bbox="1568 550 1928 657">Condition(s)</th> </tr> </thead> <tbody> <tr> <td data-bbox="996 657 1151 911">R1</td> <td data-bbox="1151 657 1373 911">(nucleophilic substitution)</td> <td data-bbox="1373 657 1568 911">KOH Allow NaOH Ignore H₂O</td> <td data-bbox="1568 657 1928 911">aqueous and heat Allow aqueous ethanol for aqueous Allow warm/reflux for heat</td> </tr> <tr> <td data-bbox="996 911 1151 1070">R2</td> <td data-bbox="1151 911 1373 1070">elimination</td> <td data-bbox="1373 911 1568 1070">KOH Allow NaOH</td> <td data-bbox="1568 911 1928 1070">(heat in ethanol)</td> </tr> <tr> <td data-bbox="996 1070 1151 1225">R3</td> <td data-bbox="1151 1070 1373 1225">nucleophilic substitution</td> <td data-bbox="1373 1070 1568 1225">(NH₃)</td> <td data-bbox="1568 1070 1928 1225">alcohol/ethanol and heat/warm and under pressure/sealed</td> </tr> <tr> <td data-bbox="996 1225 1151 1401">R4</td> <td data-bbox="1151 1225 1373 1401">nucleophilic substitution</td> <td data-bbox="1373 1225 1568 1401">KCN Allow NaCN Ignore HCN</td> <td data-bbox="1568 1225 1928 1401">(heat in ethanol)</td> </tr> </tbody> </table>	Reaction	Name of mechanism	Formula of Reagent	Condition(s)	R1	(nucleophilic substitution)	KOH Allow NaOH Ignore H ₂ O	aqueous and heat Allow aqueous ethanol for aqueous Allow warm/reflux for heat	R2	elimination	KOH Allow NaOH	(heat in ethanol)	R3	nucleophilic substitution	(NH ₃)	alcohol/ethanol and heat/warm and under pressure/sealed	R4	nucleophilic substitution	KCN Allow NaCN Ignore HCN	(heat in ethanol)	(4)
Reaction	Name of mechanism	Formula of Reagent	Condition(s)																				
R1	(nucleophilic substitution)	KOH Allow NaOH Ignore H ₂ O	aqueous and heat Allow aqueous ethanol for aqueous Allow warm/reflux for heat																				
R2	elimination	KOH Allow NaOH	(heat in ethanol)																				
R3	nucleophilic substitution	(NH ₃)	alcohol/ethanol and heat/warm and under pressure/sealed																				
R4	nucleophilic substitution	KCN Allow NaCN Ignore HCN	(heat in ethanol)																				

Question Number	Answer	Additional Guidance	Mark
23(c)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • any two or three correct scores (1) • any four or five correct scores (2) • any six or seven correct scores (3) • all eight correct scores (4) <p>P1: lone pair on N of left hand NH₃</p> <p>P2: curly arrow from NH₃ lone pair to C^(δ+)</p> <p>P3: correct dipole on C^{δ+}-Br^{δ-} bond</p> <p>P4: curly arrow from C-Br bond to Br^{δ-}</p> <p>P5: positive charge on N of ammonium salt</p> <p>P6: lone pair on N of right hand NH₃</p> <p>P7: curly arrow from NH₃ lone pair to H of N-H</p> <p>P8: curly arrow from N-H bond to N⁽⁺⁾</p>	<p><u>Example of completed mechanism:</u></p>  <p>Do not award incorrect positioning of lone pair in P1/P6 but penalise once only</p> <p>Do not award negative charge on NH₃ in P1/P6 but penalise once only</p> <p>Do not award use of half-headed arrows in P2/P4/P7/P8 but penalise once only</p> <p>Ignore any dipole on N-H bond in P7/P8</p> <p>Do not award arrow from incorrect N-H bond in P8</p> <p>Ignore H⁺ ion / HBr / NH₄Br by-product</p>	(4)

Question Number	Answer	Additional Guidance	Mark
23(c)(iii)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li data-bbox="398 443 965 480">• (absorptions at) 2260-2215 (cm⁻¹) <li data-bbox="398 663 640 700">• (due to) C≡N 	<p>Ignore any reference to 2962-2853 (cm⁻¹) / 1485-1365 (cm⁻¹) / C-H / C-C</p> <p>M1 and M2 are standalone marks</p> <p>(1) Allow any range or number within 2260-2215</p> <p>Do not award 2260-2100 Do not award any other wavenumbers</p> <p>(1) Ignore just CN Ignore name of any functional group, even if incorrect</p> <p>Do not award C-N / C=N Do not award C≡C Do not award any other bond</p>	(2)

(Total for Question 23 = 14 marks)

Section C

Question Number	Answer	Additional Guidance	Mark
24(a)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • calculation of total reactant bond enthalpies (1) • calculation of total product bond enthalpies (1) • calculation of S–F bond enthalpy (1) 	<p><u>Example of calculation:</u></p> <p>Correct answer with some working scores (3)</p> <p>$8 \times 268 + 24 \times 151 = 2144 + 3624$ (= 5768 kJ mol⁻¹)</p> <p>$5768 - (-9672) = 5768 + 9672$ (= 15440 kJ mol⁻¹) TE on M1 (M1 value + 9672)</p> <p>$15440 \div (8 \times 6) = 321.67$ = 322 (kJ mol⁻¹)</p> <p>Ignore SF except 1SF TE dependent on use of both M1 value and 9672 Do not award just division of any number by 48</p>	(3)

Question Number	Answer	Additional Guidance	Mark
24(a)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • bond enthalpy data is for the gas phase and sulfur/S₈ is a solid • bond enthalpy is not accurate for S–S/S₈/sulfur <p>OR</p> <p>bond enthalpy is an average/mean for many compounds</p>	<p>Ignore any reference to standard conditions Ignore any reference to enthalpy measurements Ignore any reference to heat loss</p> <p>(1) Allow sulfur/S₈ is not a gas</p> <p>Do not award any reference to SF₆ and/or F₂ as solid/liquid/not a gas</p> <p>Allow bond enthalpy is an average/mean for S–S/S₈/sulfur Ignore just bond enthalpy is an average/mean Ignore any reference to F–F / S–F bond enthalpy</p> <p>(1) Allow different for many Allow molecules/bonds for compounds</p> <p>If no other mark awarded: bond enthalpy data is for gas(es) and bond enthalpy is an average/mean scores (1)</p>	(2)


Question Number	Answer	Additional Guidance	Mark
24(a)(iii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • calculation of amount of SF₆ (1) • calculation of amount of CO₂ (1) • calculation of n(CO₂) : n(SF₆) (1) 	<p><u>Example of calculation:</u></p> <p>Ignore correct use of Avogadro constant throughout</p> <p>amount = $1.00 \div 146.1$ (= $0.0068446 / 6.8446 \times 10^{-3}$) Allow use of 146 for M_r (giving $0.0068493 / 6.8493 \times 10^{-3}$)</p> <p>amount = $23900 \div 44.0$ (= 543.18)</p> <p>$543.18 \div 0.006845 = 79359$ (molecules of CO₂) TE on M1 and M2 Ignore SF Do not award $0.006845 \div 543.18 = 1.2601 \times 10^{-5}$</p>	(3)

Question Number	Answer	Additional Guidance	Mark
24(b)(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> • ClO⁻ 	<p>Allow OCl⁻</p> <p>Ignore NaClO</p> <p>Do not award HClO / any other answer</p>	(1)

Question Number	Answer	Additional Guidance	Mark
24(b)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> equilibrium 1 shifts left (to make HClO) (1) equilibrium 2 shifts to the right (to make chlorine solution) (1) chlorine gas (produced) and (which is) toxic (1) 	<p>Marks can be awarded in any order</p> <p>Allow any unambiguous indication of equilibrium 1 shifting left, eg H⁺ reacts with ClO⁻ forming HClO</p> <p>Allow any unambiguous indication of equilibrium 2 shifting right, eg HClO reacts with HCl forming Cl₂ (and H₂O)</p> <p>Accept Cl₂(g) for chlorine gas</p> <p>Allow poisonous for toxic Ignore harmful for toxic</p> <p>If no other mark awarded neutralisation / reaction of acid and alkali is (highly) exothermic scores (1)</p>	(3)

Question Number	Answer	Additional Guidance	Mark
24(b)(iii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> formula of sodium chlorate(V) product (1) rest of equation correct (1) 	<p>NaClO₃ Allow ClO₃⁻ Do not award any other chlorate product</p> <p>3NaClO → 2NaCl + NaClO₃ Allow multiples Ignore state symbols, even if incorrect</p> <p>3ClO⁻ → 2Cl⁻ + ClO₃⁻ scores (2)</p>	(2)

Question Number	Answer	Additional Guidance	Mark
24(c)	<ul style="list-style-type: none"> <li data-bbox="398 373 1111 443">• calculation of molar mass of $C_{12}H_{18}Br_6$ fire retardant (1) <li data-bbox="398 587 1111 657">• calculation of % by mass of Br to 3SF or 2SF (1) 	<p data-bbox="1144 304 1491 336"><u>Example of calculation:</u></p> <p data-bbox="1144 373 1700 405">$12 \times 12(.0) + 18 (\times 1.0) + 6 \times 79.9$</p> <p data-bbox="1144 411 1196 443">OR</p> <p data-bbox="1144 450 1375 481">$641.4 \text{ (g mol}^{-1}\text{)}$</p> <p data-bbox="1144 488 1812 520">Allow use of $A_r(\text{Br}) = 80$ giving $642 \text{ (g mol}^{-1}\text{)}$</p> <p data-bbox="1144 590 1700 622">$479.4 \div 641.4 \times 100 = 74.7 / 75(\%)$</p> <p data-bbox="1144 628 1285 660">TE on M1</p> <p data-bbox="1144 667 1756 699">Do not award TE if % by mass > 100(%)</p> <p data-bbox="1144 705 1621 737">Allow 74.8(%) from $A_r(\text{Br}) = 80$</p>	(2)

Question Number	Answer	Additional Guidance	Mark
24(d)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • y-axis label and x-axis label (1) • single distribution starting at the origin and approaching but not touching the x-axis (1) • catalyst provides (alternative route with) lower activation energy / E_a (1) • more particles/collisions have $E \geq E_a$ with catalyst (1) 	<p><u>Example of labelled Maxwell-Boltzmann distribution:</u></p>  <p>All marks are standalone</p> <p>Allow fraction / amount for number Allow molecules / atoms for particles Accept kinetic energy</p> <p>(1) Do not award two or more distributions</p> <p>(1) Allow labelled activation energies with and without catalyst on diagram</p> <p>(1) Allow annotated shaded areas on diagram Allow more particles/collisions have sufficient energy to react Ignore any reference to successful collisions</p>	(4)

(Total for Question 24 = 20 marks)

(Total for Section C = 20 marks)

(Total for Paper = 80 marks)

