

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Wednesday 20 January 2021

Morning (Time: 1 hour 20 minutes)

Paper Reference **WCH13/01**

Chemistry

International Advanced Subsidiary/Advanced Level

Unit 3: Practical Skills in Chemistry I

You must have:

Scientific calculator, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- Show all your working in calculations and include units where appropriate.

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P67747A

©2021 Pearson Education Ltd.

1/1/1




Pearson

Answer ALL questions. Write your answers in the spaces provided.

- 1 (a) A student was provided with five test tubes labelled **A**, **B**, **C**, **D** and **E**, each containing a colourless aqueous solution.

The five solutions were known to be

barium chloride

nitric acid

potassium bromide

silver nitrate

sodium carbonate

The student carried out a series of tests to identify which test tube contained which solution.

- (i) The student tested each solution using universal indicator paper. Only solution **A** turned the paper red.

Identify solution **A**.

(1)

- (ii) The student mixed 1 cm^3 of solution **A** separately with 1 cm^3 of each of the other solutions.

There was no change for three of the mixtures but effervescence was observed when solution **A** was added to solution **C**.

Identify solution **C**.

(1)

- (iii) Write an **ionic** equation for the reaction between solution **A** and solution **C**. Include state symbols.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(iv) The student then mixed 1 cm³ samples of the remaining solutions as shown in **Table 1**.

Solutions mixed	Observation
B and D	no change
B and E	cream precipitate
D and E	white precipitate

Table 1

Identify the three remaining solutions.

(3)

Solution **B**

Solution **D**

Solution **E**

(b) Three of the cations in the compounds in (a) can be identified using flame tests.

Complete **Table 2**.

(3)

Cation formula	Flame colour

Table 2

(Total for Question 1 = 10 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2 Sodium hydroxide solution reacts with carbon dioxide in the air and should be standardised before use. Ethanedioic acid may be used for this standardisation.

(a) A standard solution of ethanedioic acid, $(\text{COOH})_2$, is prepared.

- 2.40 g of solid ethanedioic acid is dissolved in approximately 100 cm^3 of deionised water in a beaker.
 - The solution is transferred into a 250.0 cm^3 volumetric flask and made up to the mark with deionised water.
- (i) Give a possible reason why any solution remaining in the beaker is washed into the volumetric flask before making up to the mark.

(1)

(ii) Calculate the concentration of this standard solution of ethanedioic acid in mol dm^{-3} .

Give your answer to an appropriate number of significant figures.

[Molar mass of ethanedioic acid = 90.0 g mol^{-1}]

(2)



- (b) A **different** standard solution of ethanedioic acid is used to determine the concentration of a sodium hydroxide solution **J**.

Procedure

Step 1 A burette is rinsed with deionised water.

Step 2 The burette is then rinsed with $0.0900 \text{ mol dm}^{-3}$ ethanedioic acid and filled with this acid solution.

Step 3 A pipette is used to transfer 25.0 cm^3 portions of solution **J** to conical flasks.

Step 4 The portions are titrated with the ethanedioic acid solution using phenolphthalein indicator.


- (i) Explain why the burette is rinsed with ethanedioic acid solution in Step 2.

(1)

- (ii) The diagram shows how the student read the filled pipette in Step 3.

Identify the **two** mistakes the student made.

(2)



(iii) The student completely emptied the pipette for each transfer in Step 3.

Explain the effect **on the titre** of completely emptying the pipette rather than leaving a small amount of solution in the tip.

(2)

(iv) State the colour **change** in the conical flask at the end-point.

(2)

From to

(c) The titration results are shown.

Titration	1	2	3
Final reading / cm ³	25.05	26.60	25.50
Initial reading / cm ³	0.00	2.00	1.00
Titre / cm ³			
Titres used in calculation of mean			


(i) Complete the table and calculate the mean titre.

(2)



(ii) Calculate the concentration of the sodium hydroxide solution in mol dm^{-3} .

The equation for the titration is



(3)

(Total for Question 2 = 15 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3 This question is about the thermal decomposition of Group 2 carbonates.

A student heated a sample of a Group 2 carbonate until no more gas was produced. The equation for the decomposition is



(a) Give a reason why the delivery tube must be removed from the water bath before removing the test tube from the heat source.

(1)

(b) The results of the experiment are shown.

Measurement	Value
Volume of carbon dioxide / cm ³	95
Mass of test tube + carbonate / g	21.69
Mass of test tube / g	21.36
Mass of carbonate / g	0.33



(i) Using the results of the experiment identify the Group 2 metal.

[Molar volume of gas at room temperature and pressure = $24.0 \text{ dm}^3 \text{ mol}^{-1}$]

(3)

(ii) The student suggested that the experiment could be made more accurate by increasing the mass of carbonate from 0.33 g to 1.00 g.

No changes to the size of the apparatus or the method of measurement of the gas produced would be made.

Comment on this suggestion.

(2)

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (c) The enthalpy change for the thermal decomposition of a carbonate, $\Delta_r H$, is difficult to measure directly.

An example of a Hess's Law cycle to determine it indirectly is



In an experiment to determine ΔH_1 , 0.050 mol of MCO_3 was placed in a 100 cm^3 beaker. 60 cm^3 of 2 mol dm^{-3} hydrochloric acid (an excess) was added and the mixture stirred. The maximum temperature rise measured was 6.0°C .

[Heat capacity of solution produced = $4.18 \text{ J g}^{-1} \text{ }^\circ\text{C}^{-1}$
Density of solution = 1.0 g cm^{-3}]

- (i) Calculate the enthalpy change, ΔH_1 , for the reaction between MCO_3 and hydrochloric acid in kJ mol^{-1} . Include a sign with your answer. (2)
- (ii) Using your answer to (c)(i), calculate the enthalpy change, $\Delta_r H$, for the thermal decomposition of this Group 2 carbonate in kJ mol^{-1} . Include a sign with your answer. (1)

(Total for Question 3 = 9 marks)



- 4 The halogenoalkane 2-chloro-2-methylpropane may be prepared from 2-methylpropan-2-ol.

Procedure

- Step 1** Add 35 cm^3 of concentrated hydrochloric acid to 8.00 g of 2-methylpropan-2-ol in a conical flask.
Swirl the mixture gently for 20 minutes.
- Step 2** Two distinct layers form. The upper (organic) layer contains the required product. The lower aqueous layer is removed using a separating funnel.
- Step 3** Add a solution of sodium hydrogencarbonate to the organic layer.
Swirl gently. Stopper the separating funnel and shake it.
Invert the separating funnel and open the tap.
- Step 4** Return the separating funnel to its upright position, remove the stopper and run off the aqueous layer. Transfer the organic layer into a clean conical flask.
- Step 5** Add some anhydrous sodium sulfate.
Leave the flask to stand and decant off the liquid.
- Step 6** Distil the liquid, collecting the product between 50°C and 52°C .
- (a) (i) The concentrated hydrochloric acid used in Step 1 was labelled



Suggest **two** safety precautions, other than wearing safety spectacles and a laboratory coat, to minimise the risk when using this reagent in Step 1.

(2)

- (ii) Explain why the product in the organic layer in Step 2 does not mix with the aqueous layer.

(2)



(iii) State why the tap of the separating funnel must be opened in Step 3.

(1)

(iv) State why anhydrous sodium sulfate is added to the organic layer in Step 5.

(1)

(v) Draw the apparatus required to distil the product and collect the distillate between 50°C and 52°C in Step 6.

(4)


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) The equation for the reaction is



The final product after distillation weighed 2.62 g.

Calculate the percentage yield.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	H	hydrogen	1
-----	----------	----------	---

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1) (2)

6.9	Li	lithium	3
9.0	Be	beryllium	4
23.0	Na	sodium	11
24.3	Mg	magnesium	12
39.1	K	potassium	19
40.1	Ca	calcium	20
85.5	Rb	rubidium	37
87.6	Sr	strontium	38
132.9	Cs	caesium	55
[223]	Fr	francium	87
[226]	Ra	radium	88
[227]	Ac*	actinium	89
138.9	La*	lanthanum	57
178.5	Hf	hafnium	72
180.9	Ta	tantalum	73
183.8	W	tungsten	74
186.2	Re	rhenium	75
190.2	Os	osmium	76
192.2	Ir	iridium	77
195.1	Pt	platinum	78
197.0	Au	gold	79
[272]	Rg	roentgenium	111
101.1	Ru	ruthenium	44
102.9	Rh	rhodium	45
106.4	Pd	palladium	46
107.9	Ag	silver	47
112.4	Cd	cadmium	48
114.8	In	indium	49
118.7	Sn	tin	50
121.8	Sb	antimony	51
127.6	Te	tellurium	52
127.7	Pb	lead	82
127.8	Tl	thallium	81
200.6	Hg	mercury	80
204.4	Po	polonium	84
207.2	Pb	lead	82
209.0	Bi	bismuth	83
209.0	Po	polonium	84
210.0	At	astatine	85
210.0	Rn	radon	86
72.6	Ge	germanium	32
74.9	As	arsenic	33
75.0	Se	selenium	34
78.9	Br	bromine	35
83.8	Kr	krypton	36
91.2	Zr	zirconium	40
91.2	Y	yttrium	39
92.9	Nb	niobium	41
95.9	Mo	molybdenum	42
95.9	Tc	technetium	[98]
98.9	Ru	ruthenium	44
101.1	Rh	rhodium	45
102.9	Pd	palladium	46
106.4	Ag	silver	47
107.9	Cd	cadmium	48
112.4	In	indium	49
114.8	Sn	tin	50
118.7	Sb	antimony	51
121.8	Te	tellurium	52
127.6	Po	polonium	84
127.7	Pb	lead	82
127.8	Tl	thallium	81
200.6	Hg	mercury	80
204.4	Po	polonium	84
207.2	Pb	lead	82
209.0	Bi	bismuth	83
209.0	Po	polonium	84
210.0	At	astatine	85
210.0	Rn	radon	86
12.0	C	carbon	6
14.0	N	nitrogen	7
16.0	O	oxygen	8
19.0	F	fluorine	9
20.2	Ne	neon	10
27.0	Al	aluminium	13
28.1	Si	silicon	14
31.0	P	phosphorus	15
32.1	S	sulfur	16
35.5	Cl	chlorine	17
39.9	Ar	argon	18
4.0	He	helium	2

Elements with atomic numbers 112-116 have been reported but not fully authenticated

* Lanthanide series

* Actinide series

140	Ce	cerium	58
141	Pr	praseodymium	59
144	Nd	neodymium	60
147	Pm	promethium	61
150	Sm	samarium	62
152	Eu	europium	63
157	Gd	gadolinium	64
159	Tb	terbium	65
163	Dy	dysprosium	66
165	Ho	holmium	67
167	Er	erbium	68
169	Tm	thulium	69
173	Yb	ytterbium	70
175	Lu	lutetium	71
232	Th	thorium	90
[231]	Pa	protactinium	91
238	U	uranium	92
[237]	Np	neptunium	93
[242]	Pu	plutonium	94
[243]	Am	americium	95
[247]	Cm	curium	96
[245]	Bk	berkelium	97
[251]	Cf	californium	98
[254]	Es	einsteinium	99
[253]	Fm	fermium	100
[256]	Md	mendelevium	101
[254]	No	nobelium	102
[257]	Lr	lawrencium	103

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

