

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

Pearson Edexcel International Advanced Level

Thursday 9 October 2025

Afternoon (Time: 1 hour 30 minutes)

Paper
reference

WCH11/01

Chemistry

International Advanced Subsidiary/Advanced Level

**UNIT 1: Structure, Bonding and Introduction to
Organic Chemistry**

You must have:

Scientific calculator

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P84360A

©2025 Pearson Education Ltd.
M:1/1/1/1/




Pearson

SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box and then mark your new answer with a cross .

1 Atomic numbers and mass numbers are used to identify isotopes.

(a) What is the mass number of the isotope ${}_{36}^{84}\text{Kr}$?

(1)

- A 36
- B 48
- C 84
- D 120

(b) Which element has the atomic number 15 and mass number 31?

(1)

- A Ga
- B P
- C Pd
- D S

(c) What is the number of electrons in the selenide ion ${}_{34}^{80}\text{Se}^{2-}$?

(1)

- A 34
- B 36
- C 80
- D 82

(Total for Question 1 = 3 marks)

Use this space for any rough working. Anything you write in this space will gain no credit.



2 Mass spectrometry is used to analyse samples of elements and molecules.

(a) What is used to accelerate ions in a mass spectrometer?

(1)

- A beam of electrons
- B electric field
- C magnetic field
- D vaporisation

(b) A sample of boron contains 20.6% ^{10}B and 79.4% ^{11}B .

What is the relative atomic mass of this sample of boron?

(1)

- A 10.21
- B 10.50
- C 10.79
- D 10.80

(c) Bromine only has the two isotopes, ^{79}Br and ^{81}Br .

A pure sample of elemental bromine, Br_2 , was analysed in a mass spectrometer.

Why did the mass spectrum have a small peak with $m/z = 80$?

(1)

- A an isotope with a mass of 80 was formed in the spectrometer
- B there was contamination in the spectrometer
- C ions can form with a 2+ charge
- D this is the mean value of the two isotopic masses

(Total for Question 2 = 3 marks)

Use this space for any rough working. Anything you write in this space will gain no credit.



- 3 An organic compound contains 53.3% carbon, 11.1% hydrogen and 35.6% oxygen by mass. The molar mass of this compound is 90.0 g mol^{-1} .


[Data: A_r H = 1.0 C = 12.0 O = 16.0]

What is the **empirical** formula of the compound?

- A $\text{C}_4\text{H}_{10}\text{O}$
 B $\text{C}_4\text{H}_{10}\text{O}_2$
 C $\text{C}_4\text{H}_5\text{O}_2$
 D $\text{C}_2\text{H}_5\text{O}$

(Total for Question 3 = 1 mark)

- 4 Strontium nitrate, $\text{Sr}(\text{NO}_3)_2$, can be prepared by the reaction of strontium carbonate with dilute nitric acid. The equation is shown.



- (a) Which is the **ionic** equation for this reaction?

(1)

- A $\text{Sr}^{2+}(\text{s}) + 2\text{NO}_3^-(\text{aq}) \rightarrow \text{Sr}(\text{NO}_3)_2(\text{aq})$
 B $\text{CO}_3^{2-}(\text{s}) + 2\text{H}^+(\text{aq}) \rightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
 C $\text{SrCO}_3(\text{s}) + 2\text{H}^+(\text{aq}) \rightarrow \text{Sr}^{2+}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
 D $\text{SrCO}_3(\text{s}) + 2\text{HNO}_3(\text{aq}) \rightarrow \text{Sr}^{2+}(\text{aq}) + 2\text{NO}_3^- + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$

- (b) What is the percentage atom economy, by mass, for this preparation of strontium nitrate?

[Data: A_r H = 1.0 C = 12.0 N = 14.0 O = 16.0 Sr = 87.6]

(1)

- A 22.7
 B 70.7
 C 77.3
 D 82.8

(Total for Question 4 = 2 marks)

Use this space for any rough working. Anything you write in this space will gain no credit.



5 Which isoelectronic ion has the largest radius?

- A Ca^{2+}
- B Cl^-
- C K^+
- D S^{2-}

(Total for Question 5 = 1 mark)

6 0.100 mol of sodium nitrate dissolves in deionised water to form 250 cm^3 of solution.

What is the solution concentration, in g dm^{-3} ?

[Data: NaNO_3 molar mass = 85.0 g mol^{-1}]

- A 0.034
- B 0.400
- C 8.50
- D 34.0

(Total for Question 6 = 1 mark)

7 Which of the giant lattices of carbon atoms contain flat hexagonal rings?

- A diamond only
- B graphene only
- C graphite only
- D graphene and graphite only

(Total for Question 7 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



8 The first ionisation energy for hydrogen is 1312 kJ mol^{-1} .

What is the ionisation energy, in **joules**, that is needed per atom of hydrogen?

[Data: Avogadro constant, $L = 6.02 \times 10^{23} \text{ mol}^{-1}$]

- A 1.09×10^{-18}
- B 2.18×10^{-18}
- C 1.09×10^{-21}
- D 2.18×10^{-21}

(Total for Question 8 = 1 mark)

9 Which molecule is polar?

- A BeCl_2
- B NCl_3
- C CCl_4
- D PCl_5

(Total for Question 9 = 1 mark)

10 Zinc has electrons in s, p and d orbitals.

How many electrons **in total** does zinc have in p orbitals?

- A 12
- B 10
- C 8
- D 6

(Total for Question 10 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



11 A sample of 6.300 g of cerium reacted completely with 1.439 g of oxygen gas.





What is the formula of the oxide formed in this reaction?

[Data: A_r Ce = 140.1 O = 16.0]

- A CeO_4
 B Ce_2O_3
 C CeO_2
 D Ce_3O_4

(Total for Question 11 = 1 mark)

12 Which compound can only be named using the *E-Z* naming system and **not** using the *cis-* and *trans-* naming system?

- A 
- B 
- C 
- D 

(Total for Question 12 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



13 Free radical substitution is the mechanism of the reaction between methane and chlorine.

(a) Which drawing of curly half-arrows shows initiation?

(1)



(b) Which equation is **not** a possible propagation step in the reaction?

(1)



(c) What differentiates a termination step from a propagation step?

(1)

- A covalent bonds are formed
- B heterolytic fission occurs
- C no free radicals are formed
- D ultraviolet radiation is not needed

(Total for Question 13 = 3 marks)

TOTAL FOR SECTION A = 20 MARKS



SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

14 Ionisation energies provide information about the orbitals in the electronic structures of atoms.

(a) State what is meant by the term 'orbital'.

(2)

.....

.....

.....

.....

(b) Explain the trend in first ionisation energies down Group 1.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(c) Explain how the changes in the first four successive ionisation energies of calcium indicate the group to which it belongs.

(2)

.....

.....

.....

.....

.....

.....

.....

(d) Write the equation for the **second** ionisation energy of fluorine. Include state symbols.

(2)

(e) The **first** ionisation energy of nitrogen is $+1402 \text{ kJ mol}^{-1}$.

Predict an approximate value for the first ionisation energy of oxygen. Justify your answer.

(3)

Value kJ mol^{-1}

Justification

.....

.....

.....

.....

.....

.....

.....



(f) The first ionisation energy of magnesium and of aluminium are shown.

Element	First ionisation energy / kJ mol^{-1}
Magnesium	+738
Aluminium	+578

Explain why the first ionisation energy of aluminium is less than that of magnesium.

(2)

.....

.....

.....

.....

(Total for Question 14 = 15 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



15 Propane and butane are used as fuels in portable heaters. They are stored under pressure in cylinders as liquids. When the cylinder valve is opened, the decrease in pressure causes the liquid fuel to change into a gas.

(a) A gas cylinder contains 18.4 dm^3 of liquid propane, C_3H_8 .

Calculate the volume of propane gas that would be produced from this quantity of liquid propane at 101 kPa and 25°C .

Include units in your answer. You **must** show your working.

Use the ideal gas equation, $pV = nRT$.

[Data: $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ Density of liquid propane = 0.585 g cm^{-3}]

(4)

(b) Gas cylinders are made of metal.

Explain, with reference to structure and bonding, why these gas cylinders can withstand the high pressures.

(2)

.....

.....

.....

.....

.....



(c) Complete combustion of butane produces more carbon dioxide than the complete combustion of propane, per mole of fuel.

- (i) Write an equation for the complete combustion of propane.
State symbols are not required.

(1)

- (ii) Determine the increase in volume, in dm^3 , of carbon dioxide gas produced per mole of butane compared to per mole of propane on complete combustion.

Use $24 \text{ dm}^3 \text{ mol}^{-1}$ for the molar volume of carbon dioxide gas.

(1)

- (iii) State an environmental problem which arises from the use of propane and butane as fuels.

(1)

.....
.....



(d) Incomplete combustion of propane produces carbon monoxide, which is a serious health hazard. The UK's Health and Safety Executive has imposed a maximum workplace exposure limit. This limit is 20 ppm of carbon monoxide, by volume, during an 8-hour period.

(i) A small workroom has a volume of 50.2 m^3 .

Calculate the **maximum** number of carbon monoxide molecules allowed in the workroom if a person is present for 8 hours.

[Data: Avogadro constant, $L = 6.02 \times 10^{23} \text{ mol}^{-1}$
Molar gas volume = $24 \text{ dm}^3 \text{ mol}^{-1}$]

(3)

(ii) Suggest **two** steps a person could take to reduce the risk associated with the hazardous carbon monoxide whilst working in this room.

(2)

.....

.....

.....

.....

.....

.....

.....

(Total for Question 15 = 14 marks)



16 This question is about organic compounds with the molecular formula C_5H_{10} .

(a) Pent-1-ene and pent-2-ene are structural isomers, but only pent-2-ene has geometric isomers.

(i) State why pent-1-ene and pent-2-ene are structural isomers.

(1)


(ii) Explain why pent-2-ene has geometric isomers.

(2)

(b) Pent-1-ene can be converted into pentane.

Add a catalyst for this conversion on the dotted line.

(1)



(c) Pent-2-ene reacts with hydrogen bromide to form two different products.

Draw the **displayed** formula of each product.

(2)

Product 1	Product 2



- (d) Both pent-1-ene and pent-2-ene react with steam in the presence of an acid catalyst to produce alcohols.

Explain, by referring to the addition of steam, how both alkenes can produce the same alcohol but can also produce different alcohols.
Include names and structures of all three alcohols.

(5)

.....

.....

.....

.....

.....

.....

.....

.....

- (e) Draw **two** repeat units of poly(pent-2-ene).


(1)



- (f) There are three branched alkenes with the molecular formula C_5H_{10} .
The skeletal formulae of two of these are shown.

Draw the **skeletal** formula of the other branched alkene of C_5H_{10} .

(1)



- (g) Five cycloalkane molecules have the molecular formula C_5H_{10} .

(i) Describe the result when bromine is added to cyclopentane in the dark.

(1)

(ii) Draw the structure of **three** of the other cycloalkane isomers of C_5H_{10} .
Do **not** draw cyclopentane.

(2)

(Total for Question 16 = 16 marks)



17 This question is about the salts ammonium sulfate and copper(II) sulfate.

(a) Ammonium sulfate has the formula $(\text{NH}_4)_2\text{SO}_4$.

Ammonium salts contain the ammonium ion NH_4^+ . This ion is formed when ammonia forms a dative covalent bond with a proton.

Explain, using electron-pair repulsion theory, how the ammonia molecule and the ammonium ion have different bond angles and shapes.

(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) Copper(II) sulfate is an ionic compound.

(i) Describe an experiment which would demonstrate the movement of the copper(II) ions in an aqueous solution of this salt.

(3)

.....

.....

.....

.....

.....


.....

.....

.....

(ii) Complete the dot-and-cross diagram of the sulfate ion.
Use dots (•) for the sulfur electrons, crosses (×) for the oxygen electrons and triangles (▲) for the extra electrons.

(2)



- (c) The double salt, $(\text{NH}_4)_2\text{Cu}(\text{SO}_4)_2$, can be made by adding ammonium sulfate solution to aqueous copper(II) sulfate. The equation for this preparation is shown.



The double salt was prepared by:

- dissolving 3.35 g of ammonium sulfate in 25 cm^3 of deionised water
- dissolving 3.96 g of copper(II) sulfate in 25 cm^3 of deionised water
- adding the two solutions together, heating and then leaving to crystallise

- (i) Calculate which of the salts was in excess.

(3)



- (ii) The crystalline salt prepared is hydrated and has the formula $(\text{NH}_4)_2\text{Cu}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$. The mass collected was 9.52 g.

Calculate the percentage yield of the double salt from this preparation.

(3)

(Total for Question 17 = 15 marks)

TOTAL FOR SECTION B = 60 MARKS
TOTAL FOR PAPER = 80 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	H	hydrogen	1
-----	----------	----------	---

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
Li	Be	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	B	C	N	O	F	He
lithium	beryllium	scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	boron	carbon	nitrogen	oxygen	fluorine	helium
3	4	21	22	23	24	25	26	27	28	29	30	5	6	7	8	9	2
23.0	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
Na	Mg	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	Al	Si	P	S	Cl	Ar
sodium	magnesium	yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	aluminium	silicon	phosphorus	sulfur	chlorine	argon
11	12	39	40	41	42	43	44	45	46	47	48	13	14	15	16	17	18
39.1	40.1	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	69.7	72.6	74.9	79.0	79.9	83.8
K	Ca	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Ga	Ge	As	Se	Br	Kr
potassium	calcium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	gallium	germanium	arsenic	selenium	bromine	krypton
19	20	57	72	73	74	75	76	77	78	79	80	31	32	33	34	35	36
85.5	87.6	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	69.7	72.6	74.9	79.0	79.9	131.3
Rb	Sr	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	In	Sn	Sb	Te	I	Xe
rubidium	strontium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	indium	tin	antimony	tellurium	iodine	xenon
37	38	57	72	73	74	75	76	77	78	79	80	49	50	51	52	53	54
132.9	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	114.8	118.7	121.8	127.6	126.9	131.3
Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Pb	Bi	Po	At	Rn	Rn
caesium	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	lead	bismuth	polonium	astatine	radon	radon
55	56	57	72	73	74	75	76	77	78	79	80	82	83	84	85	86	86
[223]	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]	[272]	204.4	207.2	209.0	[210]	[222]	[222]
Fr	Ra	Ac*	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Rg	Tl	Pb	Bi	Po	At	Rn
francium	radium	actinium	rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	roentgenium	thallium	lead	bismuth	polonium	astatine	radon
87	88	89	104	105	106	107	108	109	110	111	111	81	82	83	84	85	86

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	141	144	150	152	157	163	165	167	169	173	175
Ce	Pr	Nd	Sm	Eu	Gd	Dy	Ho	Er	Tm	Yb	Lu
cerium	praseodymium	neodymium	samarium	europium	gadolinium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
58	59	60	62	63	64	66	67	68	69	70	71
232	[231]	238	[242]	[243]	[247]	[251]	[254]	[253]	[256]	[254]	[257]
Th	Pa	U	Pu	Am	Cm	Cf	Es	Fm	Md	No	Lr
thorium	protactinium	uranium	plutonium	americium	curium	californium	einsteinium	fermium	mendeleevium	nobelium	lawrencium
90	91	92	94	95	96	98	99	100	101	102	103

* Lanthanide series

* Actinide series

